arXiv:1212.6191v1 [cond-mat.mes-hall] 26 Dec 2012

Statistical Topological Insulators

I. C. Fulga,[1] B. van Heck,[1] J. M. Edge,[1] and A. R. Akhmerov[1, 2]
[1]*Instituut-Lorentz, Universiteit Leiden, P.O. Box 9506, 2300 RA Leiden, The Netherlands*
[2]*Department of Physics, Harvard University, Cambridge, Massachusetts 02138 USA*
(Dated: December 2012)

We define a class of insulators with gapless surface states protected from localization due to the statistical properties of a disordered ensemble, namely due to the ensemble's invariance under a certain symmetry. We show that these insulators are topological, and are protected by a $\mathbb{Z}_2$ invariant. Finally, we prove that every topological insulator gives rise to an infinite number of classes of statistical topological insulators in higher dimensions. Our conclusions are confirmed by numerical simulations.

PACS numbers: 03.65.Vf, 71.20.-b, 73.20.Fz

Introduction. One common definition of a topological insulator is that it is a bulk insulator which has a gapless surface Hamiltonian that cannot be continuously transformed into a gapped one [1, 2]. The surface states of a topological insulator are protected from Anderson localization and, since there is an anomaly associated with the bulk field theory of such a material [3–5], they are also robust against interactions as long as the latter respect the discrete symmetry of the system. Other possible descriptions of topological insulators arise naturally from classifications of topological terms in nonlinear sigma-models [6], from K-theory [7], and even from string theory [8].

There are, however, several known examples of disordered systems which have a surface Hamiltonian that can be continuously deformed into a gapped one, and yet protected against Anderson localization. Remarkably, these systems are closely related to conventional topological insulators. One such example is a so-called weak topological insulator, a three-dimensional material made by stacking many layers of a two-dimensional topological insulator. Its surface has two Dirac cones which can be coupled by a mass term, producing a gapped system. Despite that, Ringel *et al.* [9] have argued that since an odd number of weak topological insulator layers is conducting, its surface must always be metallic. This prediction was tested numerically [10] and later explained by Fu and Kane [11] in terms of $\mathbb{Z}_2$ vortex fugacity of a corresponding field theory. Another example of a system avoiding localization is a topological insulator subject to a random magnetic field as long as the magnetic field is zero on average [12]. A random magnetic field opens a random sign gap in the surface dispersion, which drives the surface to a critical point of the Chalker-Coddington network model [13].

These two examples share one common trait: in order for the surface to avoid localization, the disordered ensemble has to be invariant under a certain symmetry: translation by a single unit cell in case of a weak topological insulator or time-reversal for a strong topological insulator with a random magnetic field. Another common feature is that the surface theories of both of these systems are in the middle of a topological phase transition. We show that these two features define a broad class of systems, which we call statistical topological insulators (STI). An STI is an ensemble of disordered systems invariant under a certain average symmetry, in which each system may have discrete symmetries. The elements of this ensemble have surfaces protected from localization due to the combined presence of the average symmetry and the discrete ones, if any. For example, in the case of a weak topological insulator the average symmetry is translation by one unit cell, while the discrete symmetry is time-reversal.

Some STIs without disorder become topological crystalline insulators, introduced by Liang Fu [14, 15], or in other words, they have a gapless surface dispersion protected by their crystalline symmetry. Nevertheless, we show that not all the topological crystalline insulators become STIs once disorder is added, and that the ensemble symmetry need not be crystalline, as in the case of a topological insulator in a random magnetic field. We also show that STIs are a true bulk phase: in order for the surface to become localized without breaking the symmetries, the bulk must undergo a phase transition. This significantly extends the list of possible topological phases: since the bulk transition of an STI is a topological phase transition by itself, it is possible to construct a higher dimensional system with its surface pinned to the middle of an STI phase transition. Such a construction makes every single topological phase transition [6, 7] give rise to infinitely many higher-dimensional descendant topological phases.

In the following, we start by defining the STI topological invariant when the average symmetry group that protects it is $\mathbb{Z}_2$. Then we show how to build a tight-binding model for an STI in any dimension and symmetry class. Finally we illustrate the properties of STIs numerically in three 2D examples.

Construction of an STI topological invariant. To determine the necessary conditions required to obtain a statistical topological insulator, let us consider an ensemble of d-dimensional systems which are semi-infinite along



FIG. 1. (a), (b): Examples of possible patterns of surface gap closing as a function of the parameter λ, interpolating between a system and its symmetry reverse, and the surface momenta $\boldsymbol{k}$. Each gap closing contributes ±1 to the total change of the topological invariant. A nontrivial STI (a) has an odd number of gap closings at $\lambda = 0$, while a trivial one (b) has this number even, such that the total change of topological invariant from $\lambda = 1$ to $\lambda = -1$ may be 0. (c): the pattern of gap closings of an STI with two independent ensemble symmetries. Since there is an odd number of gap closings that map on themselves under both symmetries, the change of a $d-1$-dimensional STI invariant $\mathcal{Q}$ has to be odd.

at least one spatial direction, such that they have one $(d-1)$-dimensional surface. We require that the Hamiltonian H_i of each element of the ensemble is local and the correlation function of the matrix elements of this Hamiltonian is sufficiently short range. Additionally, we require that the bulk is insulating, while the surface may have surface states. We consider $d \geq 2$, so that both the surface and the bulk are self-averaging [16]. Let every ensemble element H_i be equally likely to appear as another one, $\pm\mathcal{U}H_i\mathcal{U}^{-1}$, obtained from H_i by applying a unitary or an anti-unitary $\mathbb{Z}_2$ symmetry $\mathcal{U}$. Examples of such a symmetry are mirror, time-reversal, inversion, particle-hole, and chiral symmetries.

Every element of the ensemble has the same discrete symmetries, allowing its surface to have a given topological invariant Q_{d-1}. We consider elements of an ensemble of finite but large systems of size L closed with twisted periodic boundary conditions in all directions apart from the one perpendicular to the surface. In this way we may study $d-1$-dimensional bands formed on the surface. In $d \geq 2$, the surface is macroscopic, so an insulating phase would have a definite value of Q_{d-1} in the limit $L \to \infty$.

We may calculate the amount by which Q_{d-1} changes under application of $\mathcal{U}$. Since the d-dimensional bulk is gapped, this change equals the sum of the changes of Q_{d-1} with each closing of the surface gap through a continuous path which connects H_i and $\pm\mathcal{U}H_i\mathcal{U}^{-1}$. In order to make use of the average symmetry we choose this path $H(\lambda)$ such that $H(1) = H_i$ and $H(\lambda) = \pm\mathcal{U}H(-\lambda)\mathcal{U}^{-1}$. The contribution of each gap closing to the topological invariant can be evaluated using the Green's function formalism [18–20], which allows to evaluate the topological invariant change Q_{d-1} from the surface Green's function as long as the bulk stays gapped. Each gap closing contributes a change of ±1 to the topological invariant.

All the gap closings belong to one of two groups: they either map on themselves under $\mathcal{U}$, or on another gap closing, as shown in Fig. 1a). The total change of the topological invariant can be expressed as a sum over these two types of gap closings,

$$\begin{aligned}\Delta Q_{d-1} &\equiv Q_{d-1}(\lambda=+1) - Q_{d-1}(\lambda=-1)\\ &= 2\sum_{n:\lambda_n<0} q_n + \sum_{n:\lambda_n=0} q_n,\end{aligned} \tag{1}$$

where λ_n is the value of λ where n-th gap closing is located and $q_n = \pm 1$ is its contribution to ΔQ_{d-1}. The parity of ΔQ_{d-1} equals to the parity of the last term of Eq. (1). This parity should be the same for any element of the ensemble, since a path $H_j \to H_i \to \pm\mathcal{U}H_i\mathcal{U}^{-1} \to \pm\mathcal{U}H_j\mathcal{U}^{-1}$ has the same number of gap closings at the symmetric point as the path $H_i \to \pm\mathcal{U}H_i\mathcal{U}^{-1}$. Furthermore, this parity cannot be changed by perturbations that do not close the bulk gap, since the gap closings carry a finite topological invariant, and cannot be removed by local perturbations. A single gap closing can also not be shifted away from $\lambda = 0$ since this would break the symmetry.

Let us consider the situation when ΔQ_{d-1} is odd. This means that H_i and $\pm\mathcal{U}H_i\mathcal{U}^{-1}$ may not have the same topological invariant. Since these are equally probable elements of the ensemble and Q_{d-1} is a self-averaging quantity in insulating phases, this means that the surface may not be in an insulating phase. On the other hand, in the case when ΔQ_{d-1} is even, as shown in Fig. 1b), the ensemble symmetry does not prevent the trivial insulating phase with $Q = 0$ from appearing, and does not protect the surface states from localization. We thus define $\mathcal{Q} = (-1)^{\Delta Q_{d-1}}$ as the $\mathbb{Z}_2$ topological invariant of an STI. The STI topological invariant $\mathcal{Q}$ is a bulk property, e.g. the parity of the mirror Chern number, the weak or strong topological insulator invariant. Nevertheless, an explicit evaluation of $\mathcal{Q}$ for large disorder strength is in general a hard problem, since it is necessary to find a Hamiltonian with exact symmetry $\mathcal{U}$, which can be connected to an element of the ensemble without closing the bulk gap.

The same construction works if we use an STI topological invariant $\mathcal{Q}$ as the surface topological invariant Q_{d-1}. We now consider a system with two independent ensemble $\mathbb{Z}_2$ symmetries $\mathcal{U}_1, \mathcal{U}_2$. Let $\mathcal{Q}$ be the parity of the changes of a topological invariant of a $d-2$-dimensional edge of the surface under application of $\mathcal{U}_1$. We will now calculate the change of $\mathcal{Q}$ under applying a different symmetry $\mathcal{U}_2$. Now, for each element of the ensemble H_i we need to consider a family of Hamiltonians $H_i(\lambda_1, \lambda_2)$, such that $H_i(\lambda_1, \lambda_2) = \pm\mathcal{U}_1 H(-\lambda_1, \lambda_2)\mathcal{U}_1^{-1}$ and $H_i(\lambda_1, \lambda_2) = \pm\mathcal{U}_2 H(\lambda_1, -\lambda_2)\mathcal{U}_2^{-1}$. Using dimensional reduction arguments of Ref. [20], we obtain that the parity of $\Delta\mathcal{Q}$ equals the parity of the total topological invariant carried by the gap closings inside the square

$|\lambda_1|, |\lambda_2| < 1$. Consequently, $\mathcal{Q}$ should change under applying $\mathcal{U}_2$ if and only if the number of gap closings that are mapped on themselves under application of $\mathcal{U}_1$ as well as $\mathcal{U}_2$ is odd, as shown in Fig. 1c). Repeating the same construction allows to construct STIs in dimension d using an ensemble $\mathbb{Z}_2^n$ symmetry and a $d-n$ topological invariant.

STIs with mirror symmetry. Ensemble mirror symmetry allows to construct STIs using any d-dimensional topological invariant Q. Consider an infinite set of d-dimensional layers stacked in the $d+1$-st dimension and weakly coupled to each other. We alternate layers of A- and B-type characterized by bulk topological invariants Q_A and Q_B respectively. We take Q_B to be the inverse group element of Q_A (i.e. $Q_B = -Q_A$ for $\mathbb{Z}$ invariants and $Q_B = Q_A$ for $\mathbb{Z}_2$), so that coupling two such layers results in a topologically trivial system. Such a model has a reflection symmetry with respect to each layer. Once an onsite disorder with no long range correlations is added, this symmetry is broken, and becomes an ensemble symmetry instead.

For a sufficiently weak disorder we may choose the clean system as $H(\lambda = 0)$. Let us now evaluate the total number of the fermi surfaces at the symmetric point. The general form of the Bloch wave function for this system is $\Psi(x) = \exp(ikx)\,(\Psi_A, \Psi_B)$, with Ψ_A and Ψ_B the wave function on A and B layers respectively, x the number of the unit cell, and k Bloch momentum. At $k = \pi$ the eigenstates of the Hamiltonian should also be eigenstates of mirror symmetry. These are $\exp(i\pi x)(\Psi_A, 0)$ and $\exp(i\pi x)(0, \Psi_B)$, so they belong exclusively to either A- or B-type layers. They also must carry a total topological invariant Q_A and Q_B at zero excitation energy, since the layers are characterized by a non-zero bulk topological invariant. At a finite value of λ the eigenstates of mirror symmetry couple, such that according to the dimensional reduction arguments of Ref. [20] the total topological invariant of the gap closings that map on themselves under the symmetry is Q_A. Hence if each layer carries an odd number of topologically protected edge states, the layered system is a nontrivial STI and has $\mathcal{Q} = -1$.

Numerical Simulations. As an example of a two-dimensional STI obtained via a procedure similar to the one described above, we consider a stack of coupled Kitaev chains oriented in the y-direction. The two dimensional lattice Hamiltonian of this system has the form

$$H = (2t\cos k_y - V)\sigma_z + \Delta\sigma_y \sin k_y + \alpha\sigma_x \sin k_x, \quad (2)$$

where σ_i are Pauli matrices in Nambu space, and x, y are integer coordinates perpendicular and along the chain direction. The Hamiltonian parameters are chosen as follows: the normal hopping is $t = 1$, the onsite disorder potential V is uniformly distributed in the interval $[-\varepsilon/2, \varepsilon/2]$, the p-wave pairing strength within each chain is $\Delta = 1$, and the interlayer coupling strength is $\alpha = 0.45$. This Hamiltonian belongs to symmetry class D with particle-hole symmetry $H = -\sigma_x H^* \sigma_x$. It is also related to a Hamiltonian with an ensemble mirror symmetry by a gauge transformation $\psi(x) \to (-1)^{\lfloor x/2 \rfloor}\psi(x)$.



FIG. 2. Transmission through a stack of 49 coupled Kitaev chains, each chain 150 sites long with hard wall (green) and periodic (red) boundary conditions as a function of disorder strength ε. Each point is obtained by averaging over a number of approximately 6×10^3 different disorder realizations, with error bars smaller than symbol sizes. The inset shows the sample geometry with Kitaev chains drawn as black lines, and with leads attached shown as gray rectangles.

The transmission in the x-direction of a stack of such Kitaev chains with periodic and hard wall boundary conditions in y-direction is shown in Fig. 2. The clean system with $\varepsilon = 0$ and hard wall boundary conditions has transmission $T = 2$ due to a non-chiral Majorana mode being present at each edge. Despite the fact that disorder reduces this transmission, it only disappears after the bulk goes through a delocalization transition to become a trivial Anderson insulator at a sufficiently high disorder strength.

To test the properties of the transmitting edges, we calculate the dependence of the transmission through the edges on the number of Kitaev chains N at a fixed ε. In order to verify that it is the parity of the number of gap closings that determines whether an ensemble is an STI or not, we compare this result to the same calculation done for a mirror symmetric stack of chains in symmetry class BDI with alternating topological invariants $Q = \pm 2$. Our results are shown in Fig. 3. We find that $T \sim N^{-1/2}$ for the stack of Kitaev chains, as expected for a chain of randomly coupled Majorana fermions, or more generally for 1D systems at the critical point [22–25]. Also in agreement with our expectations, we find that the edges of a BDI stack are localized with $T \sim \exp(-cN)$.

To test STIs in a dimension two higher than the dimension of the original topological invariant, we consider a triangular lattice of Majorana fermions [26–28], which is a surface model of a 3D array of coupled Kitaev chains.



FIG. 3. Transmission as a function of number of Kitaev chains (blue, solid line) or of number of BDI chains with $Q = \pm 2$ (red, dashed line) in a stack. The transmission through the Kitaev chain stack scales as $1/\sqrt{N}$, while transmission through the BDI stack is exponentially decaying. The inset shows the same curves in log-linear scale. In both cases, each point was obtained by averaging approximately 2×10^4 disorder realizations at a fixed disorder strength $\varepsilon = 3t$ and chain length of 40 lattice sites.



FIG. 4. *Top left*: Arrows show the directions in a triangular Majorana lattice in which we choose the average hopping values to be positive. *Bottom left*: The setup in which we calculate transmission through disordered Majorana lattice. The leads are shown as grey rectangles. *Right*: Transmission through the Majorana lattice as a function of system size. The red curve corresponds to the case of hoppings t_{ij} uniformly distributed in a range $[-t, 5t]$. The other three curves correspond to log-normal distributed t_{ij}, with zero log-mean and different values of the log-variance σ.

The tight binding Hamiltonian of this lattice is given by

$$H = \sum_{\langle ij \rangle} i t_{ij} \gamma_i \gamma_j, \tag{3}$$

with $t_{ij} = -t_{ji}$ real-valued couplings between Majorana fermions. In a clean translationally invariant system with one Majorana fermion per unit cell and an arbitrary choice of the hopping signs, if the hoppings have equal magnitude, then the system has a reflection symmetry with respect to a plane passing through one of the hoppings and a reflection antisymmetry (H changes sign) with respect to a perpendicular plane passing through any site. Since there is a single Fermi surface in the clean system, this means that any disorder that preserves the two mirror symmetries should make the translationally invariant triangular lattice of Majorana fermions a surface model of an STI.

We have calculated the dependence of the total transmission through a square-shaped region of a disordered Majorana lattice as a function of system size for different types of disorder. Our results are shown in Fig. 4. We have found that conductance scales positively with system size for any disorder strength both for log-normal, and uniformly distributed disorder, confirming our expectations. Our results explain the thermal metal observed in Refs. [27, 28]. These two works numerically studied a transition between two triangular Majorana lattices with Chern numbers ±1 by changing the probability distribution of hopping signs. They reported a metallic phase centered around the point where every hopping had a random sign. Since in that case the ensemble has a sufficiently high symmetry, and in particular since it has the mirror symmetries described in the previous paragraph, the metallic phase is a consequence of it being a surface model of an STI.

Conclusions and discussion. To conclude, we have unified several known examples of systems protected from localization by ensemble symmetries into the new class of statistical topological insulators. We were able to present a universal proof of why STIs avoid localization, by showing that the ensemble symmetry prevents them from having a definite value of a surface topological invariant. We have introduced a universal construction of STIs from any topological invariant using reflection symmetry, and were able to explain the thermal metal phase of Refs. [27, 28] as being a surface model of an STI.

Despite only having discussed non-interacting models, STIs should be protected from interactions, as long as the interactions do not introduce spontaneous symmetry breaking with long-range correlations. This holds since the identification of the topological invariant with a gap closing relies on the Green's function structure [20], and is thus insensitive to interactions. For weak topological insulators, the robustness against interactions was reported in Ref. [29]. A natural extension of our approach would include providing a more complete relation between ensemble symmetry groups and STIs. While we have focused on $\mathbb{Z}_2^n$ symmetries for simplicity, translational symmetries must also be sufficient to construct STI for $\mathbb{Z}_2$ topological invariant, as is the case for the weak topological insulators, or also the triangular Majorana lattice. The STIs using fractionalized phases may provide a new way to make fractional topological insulators. The way the presence of several topological insulator and STI phases in the same symmetry class enriches the phase diagram of Anderson insulators provides another interesting direction to study. Finally, we are as yet unable to solve the problem of efficiently evaluating the STI topological invariant on a general basis. It is sufficiently simple for several classes, e.g. weak TI, but becomes hard for more complicated ensemble symmetries.

Acknowledgments. The numerical calculations were performed using the KWANT package developed by A. R. Akhmerov, C. W. Groth, X. Waintal, and M. Wimmer. We would like to thank C. W. J. Beenakker, L. Fu, T. Hyart and C. R. Laumann for useful discussions. This project was supported by the Dutch Science Foundation NWO/FOM, by an ERC Advanced Investigator Grant and by the EU network NanoCTM. AA was partially supported by a Lawrence Golub Fellowship.

[1] M. Z. Hasan and C. L. Kane, Rev. Mod. Phys. **82**, 3045 (2010)
[2] X.-L. Qi and S.-C. Zhang, Rev. Mod. Phys. **83**, 1057 (2011).
[3] X.-L. Qi, T. L. Hughes and S.-C. Zhang, Phys. Rev. B **78**, 195424 (2008).
[4] S. Ryu, J. E. Moore, A. W. W. Ludwig, Phys. Rev. B **85**, 045104 (2012).
[5] Z. Wang, X.-L. Qi, and S.-C. Zhang , Phys. Rev. B **84**, 014527 (2011).
[6] S. Ryu, A. P. Schnyder, A. Furusaki, and A. W. W. Ludwig, New J. Phys. **12**, 065010 (2010).
[7] A. Yu. Kitaev, AIP Conf. Proc. **1134**, 22-30 (2009), arXiv:0901.2686.
[8] S. Ryu and T. Takayanagi, Phys. Rev. D **82**, 086014 (2010).
[9] Z. Ringel, Y. E. Kraus, and A. Stern, Phys. Rev. B **86**, 045102 (2012).
[10] R. S. K. Mong, J. H. Bardarson and J. E. Moore, Phys. Rev. Lett. **108**, 076804 (2012) .
[11] L. Fu and C. Kane, arXiv:1208.3442 (2012).
[12] K. Nomura, S. Ryu, M. Koshino, C. Mudry, A. Furusaki, Phys. Rev. Lett. **100**, 246806 (2008)
[13] J. T. Chalker and P. D. Coddington, J. Phys. C: Solid State Phys. **21**, 2665 (1988).
[14] L. Fu, Phys. Rev. Lett. **106**, 106802 (2011).
[15] T. H. Hsieh, H. Lin, J. Liu, W. Duan, A. Bansil, L. Fu, Nat. Comm. **3**, 982 (2012).
[16] It is interesting to note that although STI's do not seem possible in 1D, there is a precursor to the STI phase transition in a half-filled Anderson chain [17].
[17] M. Kappus and F. Wegner, Z. Phys. B **45**, 15 (1981).
[18] G. Volovik, *The Universe in a Helium droplet*, Oxford University Press, Oxford (2003).
[19] V. Gurarie, Phys. Rev. B **83**, 085426 (2011).
[20] A. M. Essin and V. Gurarie, Phys. Rev. B **84**, 125132 (2011).
[21] A. R. Akhmerov, J. P. Dahlhaus, F. Hassler, M. Wimmer, and C. W. J. Beenakker, Phys. Rev. Lett. **106**, 057001 (2011).
[22] P. W. Brouwer, A. Furusaki, I. A. Gruzberg, and C. Mudry, Phys. Rev. Lett. **85**, 1064 (2000).
[23] P. W. Brouwer, A. Furusaki, and C. Mudry, Phys. Rev. B 67, 014530 (2003).
[24] O. Motrunich, K. Damle, and D. A. Huse, Phys. Rev. B **63**, 224204 (2001).
[25] I. A. Gruzberg, N. Read, and S. Vishveshwara, Phys. Rev. B **71**, 245124 (2005).
[26] R. R. Biswas, L. Fu, C. R. Laumann and S. Sachdev, Phys. Rev. B **83**, 245131 (2011).
[27] Y. E. Kraus and A. Stern, New J. Phys. **13**, 105006 (2011).
[28] C. R. Laumann, A. W. W. Ludwig, D. A. Huse and S. Trebst, Phys. Rev. B **85**, 161301(R) (2012).
[29] C.-X. Liu, X.-L. Qi, S.-C. Zhang, Physica E **44**, 906-911 (2012).